

# Somatic embryogenesis and peroxidase activity of desiccation tolerant mature somatic embryos of loblolly pine

TANG Wei

(O-23, E. S. King Village, 2900 Ligon St., North Carolina State University, Raleigh, NC 27607, USA)

**Abstract:** White, translucent, glossy mucilaginous callus was initiated from the mature zygotic embryos explants on callus induction medium with 2,4-D, BA, and kinetin in the 3-9th week of culture. This type of callus induction occurred at a lower frequency with either  $\alpha$ -naphthaleneacetic acid (NAA) or IBA (both 8 mg/L). White, translucent, glossy mucilaginous callus was embryogenic and mainly developed from the cotyledons of the mature zygotic embryo. Somatic embryos were formed on differentiation medium. Desiccation tolerance can be induced by culturing somatic embryos of loblolly pine (*Pinus taeda* L.) on medium supplemented with 50  $\mu$ m abscisic acid (ABA) and/or 8.5% polyethylene glycol (PEG<sub>6000</sub>). Scanning electron microscopy of desiccated somatic embryos showed that the size and external morphology of the desiccation tolerant somatic embryos recovered to the pre-desiccation state within 24-36 h, whereas the sensitive somatic embryos did not recover and remained shriveled, after the desiccated somatic embryos had been rehydrated. Peroxidase activity of desiccated somatic embryos increased sharply after 3 days of desiccation treatment, and desiccation tolerant somatic embryos had higher peroxidase activity compared to sensitive somatic embryos. Higher peroxidase activity of desiccation tolerant somatic embryos was possibly advantage of catalyzing the reduction of H<sub>2</sub>O<sub>2</sub> which was produced by drought stress, and protecting somatic embryos from oxidative damage.

**Key words:** *Pinus taeda* L.; Somatic embryogenesis; Desiccation tolerance ; Peroxidase activity

**CLC number:** Q55; S791.255.04

**Document code:** A

**Article ID:** 1007-662X(2001)03-02147-06

## Introduction

Somatic embryogenesis has been regarded as the in vitro system of choice with the potential for eventual mass propagation of superior and genetically engineering forest tree genotypes in both coniferous and hardwood species (Gupta and Durzan 1991; Attree and Fowke 1993; Zimmerman 1993; Park *et al.* 1998). Somatic embryogenesis in conifer was first described relatively recently by three independent groups: Hakman *et al.* (1985) and Chalupa (1985) described somatic embryogenesis from cultured zygotic embryos of Norway spruce, while Nagmani & Bonga (1985) reported the induction of haploid European larch (*Larix decidua* Mill.) embryo from megagametophytic tissue. In just a few years, mature conifer somatic embryos had been produced, of a quality with compares very favourable with mature zygotic embryos. Somatic embryogenesis of loblolly pine has been induced by Gupta *et al.* 1987, Becwar *et al.* 1990, and Li *et al.* 1998, and is considered to be the most promising method for large-scale propagation of loblolly pine and it offers the potential with automated system to produce large numbers of synthetic seeds (Attree *et al.* 1993; Becwar *et al.* 1995; Gupta *et al.*

1993; Handley *et al.* 1995).

Desiccation (partial drying) can improve the germination and plant regeneration response of somatic embryos in conifers (Attree *et al.* 1993). The successful induction of desiccation tolerance in somatic embryos has been reported for some angiosperm and conifer species (Attree *et al.* 1993, Florin *et al.* 1993; Bewley 1979; Brown *et al.* 1993; Iida *et al.* 1992; Seneratna *et al.* 1990). Partial drying has been described for Skita spruce (Robertson *et al.* 1991), red spruce (Harry & Thorpe 1991), black spruce (Tremblay & Tremblay 1991), white spruce (Kong and Yeuang 1992), and Norway spruce (Bozhkov *et al.* 1992). Robertson *et al.* (1990) first observed that partial drying of interior spruce somatic embryos at high relative humidity (RH>95%) increased germination frequency, decreased germination times, and improved the synchrony of root and shoot elongation compared to untreated somatic embryos. Of various chemicals and physical stresses have been applied to induce desiccation tolerance, abscisic acid (ABA) and polyethylene glycol (PEG) have been frequently employed (Attree *et al.* 1993; Florin *et al.* 1993; Zeevart *et al.* 1988). ABA and PEG influence the accumulation of specific mRNAs, storage reserves, storage lipid triacylglycerol (TAG) in conifer somatic embryos and zygotic embryos (Attree *et al.* 1992), of which some may be involved in seed protection. In some plant species, late embryogenesis abundant (LEA) proteins, whose accumulation increase dramatically at the beginning of zygotic embryo desiccation and which are regulated by ABA, are considered to be involved in protecting tissue from desiccation (Dure *et al.* 1993). Leal

<sup>1</sup> **Biography:** TANG Wei (1964-), Research associate, O-23, E. S. King Village, 2900 Ligon St., North Carolina State University, Raleigh, NC 27607, USA. E-mail: wtang638@yahoo.com

**Received date:** 2001-02-26

**Responsible editor:** Chai Ruihai

and Misra (1993) reported that late developmental stage of white spruce zygotic embryos accumulated high levels of LEA transcripts. However, little is known about precise mechanism by which ABA and PEG induce desiccation tolerance in conifer somatic embryos. Peroxidase catalyzes the reduction of  $\text{H}_2\text{O}_2$  and protects embryos from oxidative damage. There is no report about peroxidase activity of desiccation tolerant somatic embryos in conifer. In this investigation, we report the first evidence of increased peroxidase activity in desiccation tolerant somatic embryos of loblolly pine.

## Materials and Methods

### Plant material

Mature cones of genotypes E-822, E-311, and E-440 of loblolly pine were collected from 25-30 year old trees in the morning hours in October, 1996, from Yingde Seed Orchard at Yingde, Guangdong Province, and Shaoyang Seed Orchard at Shaoyang, Hunan Province, China. All cones were stored in plastic bags at 4 °C for 2 - 18 months. Seeds were disinfected by immersion in 70% v/v ethanol for 30 s and in 0.1% mercuric chloride for 20 min, followed by 4-5 rinses in sterile distilled water. Mature zygotic embryos were aseptically removed from the seeds and placed horizontally on a solidified callus induction medium in 100 mL Erlenmeyer flasks (40 mL medium in 100 mL flask, 5-8 explants per flask).

### Callus and somatic embryo induction

Callus and somatic embryo induction were carried out as described previously (Tang *et al.* 1998a). Mature zygotic embryos were cultured on callus induction medium consisted of TE (Tang *et al.* 1998) supplemented with 8 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D), 4 mg/L 6-benzylaminopurine (BA), and 4 mg/L Kinetin (KT). Callus was formed on cotyledons, hypocotyls, and radicles of mature zygotic embryos after 9 weeks on induction medium. Then callus was subcultured on the same medium with lowered 1/5 auxin and cytokinin concentrations. After 9 weeks, white, translucent, glossy mucilaginous calluses containing embryogenic suspensor masses (ESM) and immature somatic embryos was selected for proliferation on the medium consisted of TE medium supplemented with 2 mg/L 2,4-D, 0.5 mg/L BA, 0.5 mg/L indole-butyric acid (IBA), and 0.5 mg/L Kinetin. Then the influences of different concentrations of ABA, PEG<sub>6000</sub>, and activated charcoal were tested. Each treatment was comprised of 150-200 explants and the experiment was repeated thrice. In the callus induction and proliferation media, the concentrations of sucrose, casein hydrolysate, glutamine, myo-inositol, and agar were 30 g/L, 500 mg/L, 500 mg/L, 500 mg/L, and 7 g/L, respectively. The pH was adjusted to 5.8 with 1N KOH or 0.5 N HCl prior to autoclaving at 121°C for 18 min. Callus induction took place in the darkness at 23°C. Data were analyzed with Analysis of Variance, and mean comparison

was made with the Least Significant Difference test at 5% level of probability.

### ABA and PEG treatment, desiccation, and rehydration

After 9 weeks of culture, mature somatic embryos were treated with ABA and PEG<sub>6000</sub> to induce desiccation tolerance. The culture medium was replaced by TE supplemented with 50  $\mu\text{M}$  ABA and/or 8.5% PEG<sub>6000</sub>. As a control desiccation-sensitive preparation, the medium was replaced by TE without ABA and PEG<sub>6000</sub>. After 7 days of incubation in dark at 25°C, embryos were selected by sieving through a 200  $\mu\text{m}$  nylon mesh and then washed with sterile deionized water. These consisted of cotyledon-stage embryos. They were transferred to a sterile filter paper, which was moistened with a few drops of sterile deionized water, in a 60 mm x 15 mm plastic Petri dish. Somatic embryo desiccation was performed according to the method of Takahata *et al.* (1993). Embryos in a Petri dish were dried through a series of desiccators in which the relative humidity (RH) was kept constant using a saturated solution of  $\text{K}_2\text{SO}_4$  (RH 87%),  $\text{Na}_2\text{CO}_3$  (RH 80%),  $\text{NaCl}$  (RH 70%),  $\text{NH}_4\text{NO}_3$  (RH 61%),  $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$  (RH 50%) or  $\text{K}_2\text{CO}_3 \cdot 1.5\text{H}_2\text{O}$  (RH 40%). They were transferred daily from a desiccator at a higher RH to one at a lower RH. After the desiccation treatment, the filter paper bearing the embryos was transferred to TE agar (0.7%)-solidified medium containing 2% sucrose and incubated at 25°C under a 16-h photoperiod with light provided by cool white fluorescent bulbs ( $75 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ).

### Scanning electron microscopy

Embryogenic calluses were prepared for scanning electron microscopy according to Fowke *et al.* (1994) with some modification. Tissues were fixed overnight in 4% glutaraldehyde and 100 mM phosphate buffer (pH 7.0), washed one time in 100 mM phosphate buffer (pH 7.0) for 30 min, followed by dehydration in successive ethanol alcohol solution of 85%, 95%, and 100%, each repeated twice for 5 min. Specimens were dried in a critical-point-drier with  $\text{CO}_2$  for 2 h, mounted on Cu stubs and gold-coated. The samples were examined and photographed in HITACHI S-800 scanning electron microscope.


### Peroxidase activity

The enzyme activity was assayed according to the procedure reported by Kim and Yoo (1996), 0.1 g FW somatic embryos (30-50 somatic embryos) was homogenized in 1 mL phosphate buffer (pH 6.0, 0.1M) for 5 min. Homogenates were then centrifuged at 12 000 rpm for 5 min at 4°C and the supernatant was used for peroxidase activity assay. Peroxidase activity was determined at 30°C with a spectrophotometer (470 nm) following the formation of tetraguaiacol in a 3-mL reaction mixture containing 1 mL of 0.1 M phosphate buffer (pH 6.0), 1 mL of 15 mM 2-methoxyphenol (guaiacol); 1 mL of 3 mM  $\text{H}_2\text{O}_2$ ; and 5  $\mu\text{L}$  of enzyme extract. One unit of peroxidase activity (U)

represents the amount of enzyme catalyzing the oxidation of 1 mol of guaiacol in 1 min. Data were analyzed by an Analysis of Variance, and mean comparisons were made with the Least Significant Difference test at 5% level of probability.

## Results and Discussion

Callus induction and somatic embryos maturation. Calli were formed from the mature zygotic embryos explants on media containing auxin and cytokinin in the 3-9th week of culture. As the calli started to proliferate, four types could be distinguished: white, translucent, glossy mucilaginous, light yellowish, loose, glossy globular, light green globular, and light brown globular. White, translucent, glossy mucilaginous callus was initiated from 16.7%, 19.8%, and 11.7% of the E-822, E-311, and E-440 mature zygotic embryos, respectively, on TE callus induction medium with 2,4-D, BA, and kinetin (Fig. 1). This type of callus induction occurred at a lower frequency with either  $\alpha$ -naphthaleneacetic acid (NAA) or IBA (both 8 mg/l). White, translucent, glossy mucilaginous callus developed from the cotyledons of the mature zygotic embryo, light yellowish, loose, glossy globular, light green globular, and light brown globular calli formed from the hypocotyl as well as from the radicle of the explants. White, translucent, glossy mucilaginous callus was embryogenic, and was selectively proliferated on solid callus proliferation medium with 2 mg/L 2,4-D, 0.5 mg/L BA, 0.5 mg/L indole-butyric acid (IBA), and 0.5 mg/L kinetin or liquid callus proliferation medium supplemented with the same plant growth regulators as solid medium.




**Fig. 1 Genotypes and the time of callus induction on the induction frequency of embryogenic callus cultures derived from mature zygotic embryos of loblolly pine**

Each treatment was replicated three times, and each replicate consisted of 30–50 somatic embryos. Values represent the means  $\pm$  S.D.

In order to encourage the maturation of loblolly pine somatic embryos, embryogenic tissue cultures containing embryogenic suspensor masses, immature somatic em-

bryos, and globular somatic embryos must be transferred from an environment that promotes cleavage polyembryogenesis to one containing ABA and ideally raised osmotic concentration. ABA, PEG, and activated charcoal are important for the maturation of loblolly pine somatic embryos. ABA prevents the developing embryos from germinating precociously; PEG encourages the maturation of somatic embryos by regulating the osmotic pressure. Activated charcoal absorbs harmful compounds and hormones in tissues and encourages the maturation of somatic embryos (Attree *et al.* 1993; Gupta *et al.* 1993; Tautoris *et al.* 1991). However, the optimal ABA, PEG, and activated charcoal concentration required to promote maturation of loblolly pine somatic embryos needs to be determined. The highest frequency of somatic embryos (both precotyledonary somatic embryos and cotyledonary somatic embryos per gram cultures with the period of 12 weeks can be obtained when ABA, PEG<sub>6000</sub>, and activated charcoal concentration are 4mg/L, 75g/L, and 5g/L, respectively (Fig. 2).




**Fig. 2 Influence of ABA, PEG<sub>6000</sub>, and activated charcoal on the number of somatic embryos per gram embryogenic callus cultures derived from mature zygotic embryos of loblolly pine.** I: 2 mg/L ABA, 20 g/L PEG<sub>6000</sub>, and 2 g/L activated charcoal; II: 3 mg/L ABA, 50 g/L PEG<sub>6000</sub>, and 3 g/L activated charcoal; III: 4mg/L ABA, 75 g/L PEG<sub>6000</sub>, and 5 g/L activated charcoal; IV: 5mg/L ABA, 100 g/L PEG<sub>6000</sub>, and 7 g/L activated charcoal. Each treatment was replicated three times, and each replicate consisted of 30–50 somatic embryos. Values represent the means  $\pm$  S.D.

Scanning electron microscopy of desiccation tolerant somatic embryos. This investigation showed that globular somatic embryos may be obtained from embryogenic calluses cultured on callus proliferation medium, but cotyledonary somatic embryos can be produced only from embryogenic calluses cultured on medium supplemented with ABA and PEG<sub>6000</sub> (Fig. 3a-d). No visual differences in morphological appearance were observed between the somatic embryos treated and untreated with ABA and PEG<sub>6000</sub>, except that the addition of ABA and PEG<sub>6000</sub> reduced embryo growth slightly. Scanning electron micro-

scopical observation of non-desiccation embryos revealed that their epidermal cells were arranged regularly and that their cell surface was smooth, similar to the appearance reported by Nitta *et al.* (1997) and Wakui *et al.* (1999). Scanning electron microscopy of desiccated somatic embryos revealed that the external morphology of the desiccation tolerant and sensitive embryos differed. The whole surface of the desiccation tolerant embryos showed that a mesh like structure due to wrinkling of the cells, whereas the surface of desiccation sensitive embryos appeared to be torn and less wrinkled than those of the desiccation tolerant embryos. A similar morphology was reported in

desiccation tolerant somatic embryos of white spruce (Fowke *et al.* 1994) and Brassica napus L. (Wakui *et al.* 1999). After transfer to TE germination medium (Tang *et al.* 1998a), desiccation sensitive embryos did not show recovery of their morphology to that before desiccation, and reminded shriveled with an irregular cell surface after 7-10 days of rehydration (Fig 3e). Desiccation tolerant somatic embryos imbibed water rapidly and within 24-36 h had regained the same size and appearance as they had before desiccation. Their epidermal cells appeared turgid and were arranged regularly (Fig. 3f). Most of these somatic embryos develop roots within 7-10 days.



**Fig. 3 Scanning electron microscopy of somatic embryogenesis in loblolly pine**

a--Embryogenic callus (Bar = 1cm), b--Embryogenic suspensor mass (Bar = 0.1cm), c--Globular somatic embryo (Bar = 0.1mm), d--Cotyledonary somatic embryo (Bar = 0.2mm), e--Desiccation sensitive somatic embryo with an irregular cell surface (Bar = 0.3mm), f--Desiccation tolerant somatic embryo with regularly arranged epidermal cells (Bar = 0.3mm).

Peroxidase activity of desiccation tolerant somatic embryos.  $H_2O_2$  levels increase in plant cells after exposure to many environmental stresses such as radiation, drought, wounding, extreme temperature, and pathogen attack (Baker and Orlandi 1995; Yahraus *et al.* 1995). Prolonged accumulation of toxic amount of  $H_2O_2$  within the cell is eventually lethal. To avoid these deleterious effects, all organisms express peroxidase, which function as detoxifying enzymes by catalyzing the reduction of  $H_2O_2$  (Creissen *et al.* 1994). This investigation results indicated that desiccation tolerant somatic embryos induced by ABA and/or PEG have higher peroxidase activity, which is advantage of the reduction of  $H_2O_2$  produced by desiccation and the survival of the somatic embryos, as well as the maintenance of the tissue system during dehydration, protecting

embryos from oxidative damage, compared to desiccation sensitive somatic embryos. Peroxidase activity of desiccation tolerant somatic embryos increased sharply after 3 days of desiccation treatments (Table 1). It has been shown that ABA affects the stabilization of the cell membrane system (Gadea *et al.* 1999), induces the expression of late embryogenesis abundant (LEA) proteins for the stabilization of cell structure under dehydration stress (Dure 1993), and stimulates the accumulation of the storage lipid triacylglycerol (TAG) (Attree *et al.* 1992). This investigation suggested that ABA and/or PEG<sub>6000</sub> also probably promoted the production of peroxidase of desiccation tolerant somatic embryos, which catalyzes the reduction of  $H_2O_2$  and protect embryos from oxidative damage.

**Table 1 Peroxidase activity of desiccation tolerant mature somatic embryos of loblolly pine under different relative humidity (RH).**

Treatment	Peroxidase activity (U g <sup>-1</sup> FW)*					
	87%**	80%	70%	61%	50%	40%
ABA	24.3±2.1b	25.9±2.9b	30.5±2.7b	42.1±4.2b	49.7±5.1b	50.6±4.7b
PEG <sub>6000</sub>	21.5±1.6b	23.1±1.8b	27.2±1.8b	38.5±3.1b	45.2±3.5b	47.3±3.8b
ABA+PEG <sub>6000</sub>	31.7±2.9a	33.8±3.4a	36.2±3.1a	47.9±3.9a	59.8±5.7a	61.2±7.6a
Control	11.8±1.3c	12.7±1.1c	13.5±1.4c	15.6±1.2c	16.2±1.1c	16.9±2.2c

\*Peroxidase activity of somatic embryos was determined after 5-8 hrs of desiccation treatment. Each treatment was replicated three times, and each replicate consisted of 30–50 somatic embryos (0.1 g FW). Values represent the means ± S.D. Means followed by the same letter are not significantly different at the 0.05 level of confidence. \*\*Relative humidity (RH).

## Acknowledgement

This research was supported by China Postdoctoral Science Foundation (97-11). I wish to acknowledge Meibao Wu and Professor Ron Sederoff for their help to this work.

## References

- Attree, S.M., Fowke L.C. 1993. Embryogeny of gymnosperms: advances in synthetic seed technology of conifers [J]. *Plant Cell Tiss. Org. Cult.*, **35**: 1-35.
- Attree, S.M., Pomeroy M.K., Fowke L.C. 1992. Manipulation of conditions for improved triacylglycerol biosynthesis and desiccation tolerance [J]. *Planta*, **187**: 395-404.
- Baker, C.J., Orlandi, E.W. 1995. Active oxygen species in plant pathogenesis [J]. *Annu. Rev. Phytopathol.*, **33**: 299-321.
- Becwar, M.R., Nagmani R., Wann, S.R. 1990. Initiation of embryogenic cultures and somatic embryo development in loblolly pine (*Pinus taeda*) [J]. *Can. J. For. Res.*, **20**: 810-817.
- Becwar, M.R., Pullman, G.S. 1995. Somatic embryogenesis in loblolly pine (*Pinus taeda* L.) [C]. In: Jain SM, Gupta PK, Newton RJ (eds) *Somatic embryogenesis in woody plants*, vol. 3, the Netherlands: Kluwer Academic Publishers, pp 287-301.
- Bewley, J.D. 1979. Physiological aspects of desiccation tolerance [J]. *Annu. Rev. Plant Physiol.*, **30**: 195-238.
- Bozhkov, P.V., Ledebenka, L.A., Shiryayeva, G.A. 1992. A pronounced synergistic effects of abscisic acid and 6-benzyladenine on Norway spruce (*Picea abies* L.) somatic embryo maturation [J]. *Plant Cell Rep.*, **11**: 386-389.
- Brown, D.C.W, Watson, E.M., Pechan, P.M. 1993. Induction of desiccation tolerance in microspore-derived embryos of *Brassica napus* [J]. *In Vitro Cell Dev Biol* **29P**: 113-118.
- Chalupa, V. 1985. Somatic embryogenesis and plantlet regeneration from immature and mature embryos of *Picea abies* (L.) Karst [J]. *Comm. Inst. For.*, **14**: 57-63.
- Creissen, C.P., Edwards, E.A., Mullineaux, P.M. 1994. Glutathione reductase and ascorbate peroxidase [C]. In: Foyer CH, Mullineaux PM (eds) *Cause of photooxidative stress and amelioration of defense system in plants*, CRC Press, Boca Raton, pp 344-364.
- Dure LIII. 1993. Structural motifs in Lea proteins [C]. In: Close TJ, Bray EA (eds) *Plant responses to cellular dehydration during environmental stress*. The American Society of Plant Physiologists, Rockville, MD, pp.91-103.
- Florin, B., Tessereau, H., Leconteux, C., Dicher, C., Petiard, V. 1993. long-term preservation of somatic embryos [C]. In: Redenbaugh K (ed) *Synseeds*. CRC Press, Boca Raton, pp 133-162.
- Fowke, L.C., Attree, S.M., Rennie, P.J. 1994. Scanning electron microscopy of hydrated and desiccated mature somatic embryos and zygotic embryos of white spruce (*Picea glauca* [Moench] Voss) [J]. *Plant Cell Rep.*, **13**: 612-618.
- Gadea, J., Conejero, V., Vera, P. 1999. Developmental regulation of a cytosolic ascorbate peroxidase gene from tomato plants [J]. *Mol. Gen. Genet.*, **262**: 212-219.

- Gupta, P.K., Durzan, D.J. 1987. Biotechnology of somatic polyembryogenesis and plantlet regeneration in loblolly pine [J]. *Bio/Technology*, **5**:147-151.
- Gupta, P.K., Durzan, D.J. 1991. Loblolly pine (*Pinus taeda* L.) [C]. In: Bajaj YPS (ed) *Biotechnology in agriculture and forestry*, vol 16, Trees III, Berlin: Springer-Verlag, pp 383-407.
- Gupta, P.K., Pullman, G., Timmis, R., Kreitinger, M., Carlson, W.C., Grob, J., Welty, E. 1993. Forestry in the 21st Century: The biotechnology of somatic embryogenesis [J]. *Bio/Technology*, **11**: 454-459.
- Hakman, I., von Arnold S. 1985. Plant regeneration through somatic embryogenesis in *Picea abies* (norway spruce) [J]. *J Plant Physiol.*, **121**:49-158.
- Handley L.W., Becwar, M.R., Chesick, EE, 1995. Research and development of commercial tissue culture system in loblolly pine [J]. *Tappi J.*, **78**(5): 169-175.
- Harry, I.S., Thrope, T.A. 1991. Somatic embryogenesis and plant regeneration from mature zygotic embryos of red spruce [J]. *Bot Gaz.*, **152**:446-452.
- Iida, Y., Watabe, K., Kamada, H., Harada, H. 1992. Effect of abscisic acid on the induction of desiccation tolerance in carrot somatic embryos [J]. *J Plant Physiol.*, **140**: 356-360.
- Kim, Y.H., Yoo, Y.J. 1996. Peroxidase production from carrot hairy root cell culture [J]. *Enzyme Microb Technol.*, **18**: 531-536.
- Kong, L., Yeung, E.C. 1992. Development of white spruce somatic embryos: II Continual shoot meristem development during germination [J]. *In Vitro Cell Dev. Biol.*, **28**: 125-131.
- Leal I, Misra. 1993 Developmental gene expression in conifer embryogenesis and germination. III. Analysis of crystalloid protein mRNAs and desiccation protein mRNAs in the developing embryo and megagametophyte of white spruce (*Picea glauca* (Moench) Voss) [J]. *Plant Sci.*, **88**:25-37.
- Li, X.Y., Huang, F.H, Murphy, J.B., Gbur, E.E . 1998 Polyethylene glycol and maltose enhance somatic embryo maturation in loblolly pine [J]. *In Vitro Cell Dev. Biol.*, **34**: 22-26.
- Nagmani, R., Bonga, J.M. 1985. Embryogenesis in subcultured callus of *Larix deciduas* [J]. *Can. J. For. Res.*, **15**:1088-1091.
- Nitta, T., Takahata, Y., Kaizuma, N. 1997. Scanning electron microscopy of microspore embryogenesis in *Brassica* spp. *Plant Cell Rep.*, **16**: 406-410.
- Park, Y.S., Barrett, J.D., Bonga, J.M. 1998. Application of somatic embryogenesis in high-value clonal forestry: deployment, genetic control, and stability of cryopreserved clones [J]. *In Vitro Cell Dev Biol.*, **34**: 231-239.
- Robertson, D.R., Lazaroff, W.R., Webster, F.B. 1991. Interaction between maturation and high relative humidity treatments and their effects on germination of Skita spruce somatic embryos [J]. *J Plant Physiol.*, **138**: 1-6.
- Robertson, D.R., Sutton, B.C.S, Flinn, B.S. 1990. Synchronous and high frequency germination of interior spruce somatic embryos following partial drying at high relative humidity [J]. *Can. J. Bot.*, **68**: 1086-1090.
- Seneratna, T., McKersie, B.D, Bowley S.R. 1990. Artificial seed of alfalfa (*medicago sativa* L.). Induction of desiccation tolerance in somatic embryos [J]. *In Vitro Cell Dev Biol.*, **26**: 85-90.
- Takahata, Y., brown, D.C.W., Keller, W.A., Kaizuma, N. 1993. Dry artificial seed and desiccation tolerance induction in microspore-derived embryos of broccoli. *Plant Cell Tiss Org Cult.*, **35**: 121-129.
- Tang, W., Ouyang F, Guo ZC 1998a. Studies on embryogenic callus induction and plant regeneration in loblolly pine. *Scientia Silvae Scinicae* [J], **34**: 115-119.
- Tang W, Ouyang, F., Guo, Z.C. 1998b. Plant regeneration through organogenesis from callus induced from mature zygotic embryos of loblolly pine [J]. *Plant Cell Rep.*, **17**:557-560.
- Tautorius, T.E., Fowke, L.C., Funstan, D.I. 1991. Somatic embryogenesis in conifers [J]. *Can. J. Bot.*, **69**:1873-1899.
- Tremblay, L., Tremblay, F.M . 1991. Carbohydrate requirement for the development of black spruce (*Picea mariana* (Mill)B.S.P.) and red spruce (*P. rubens* Sarg.) somatic embryos [J]. *Plant Cell Tiss Org Cult.*, **27**: 95-103.
- Wakui, K., Takahata, Y., Kaizuma, N. 1999. Scanning electron microscopy of desiccation- tolerance and - sensitive microspore-derived embryos of *Brassica napus* L. [J]. *Plant Cell Rep.*, **18**: 595-600.
- Yahraus, T., Chandra, S., Legendre, L., Lowc, P. 1995. Evidence for a mechanically induced oxidative burst [J]. *Plant Physiol.*, **109**: 1259-1266.
- Zeevart, J.A.D., Creelman, R.A. 1988. Metabolism and physiology of abscisic acid [J]. *Ann Rev Plant Physiol Plant Mol Biol.*, **39**: 439-473.
- Zimmerman, J.L. 1993. Somatic embryogenesis: a model for early development in higher plants [J]. *Plant Cell*, **5**:1411-1423.